

Byerly, W.E.

QA
211
B96

UC-NRLF



B 4 248 586

QA 211 B96

IN MEMORIAM
George Davidson
1825-1911



Professor of Geography
University of California

class
t. e.
g

Byerly, William Elmer

Math D

SYLLABUS

OF A

COURSE IN THE THEORY OF EQUATIONS.

BOSTON:

PUBLISHED BY GINN, HEATH, & CO.

1883.

COPYRIGHTED BY GINN, HEATH, & CO., 1883.

Mathematical Books.

	Intro. Price.
BYERLY: Differential Calculus	\$2.00
Integral Calculus	2.00
HALSTED: Mensuration80
HARDY: Elements of Quaternions	2.00
HILL: Geometry for Beginners80
PEIRCE: Elements of Logarithms40
Mathematical Tables, Chiefly to Four Figures25
Three- and Four-Place Tables of Logarithms40
WALDO: Multiplication and Division Tables. Folio size,	.40
Small size,	.20
WENTWORTH:	
Elements of Algebra90
Complete Algebra	1.12
Plane Geometry60
Plane and Solid Geometry	1.00
Plane Trigonometry. Paper30
Plane and Spherical Trigonometry48
Plane Trigonometry and Tables. Paper48
Plane and Spherical Trigonometry and Tables90
Five-Place Log. and Trig. Tables. Paper25
Cloth45
WENTWORTH & HILL:	
Practical Arithmetic75
WHEELER: Plane and Spherical Trigonometry80

*Copies sent to Teachers for Examination, with a view to
Introduction, on receipt of Introduction Price.*

GINN, HEATH & CO., Publishers.

BOSTON. NEW YORK. CHICAGO.

SYLLABUS

OF A

COURSE IN THE THEORY OF EQUATIONS.



1. Define a *root* of an equation. Explain a short method of substituting any given number for x in a numerical equation, using only the detached coefficients.

2. If a is a root of the equation, the first member is divisible by $x - a$. Give a short formal proof. Give a second proof showing the form of the quotient; *i.e.*, substitute a for x in the first member of the equation, and then subtract the result from the given first member, and the equation will be in a form where every term is obviously divisible by $x - a$.

3. Prove the converse of the theorem in § 2. If $x - a$ will divide the first member of the equation, a is a root.

4. Assuming that every equation has at least one root, prove that every equation of the n th degree has n roots and can be thrown into the form

$$A(x - a)(x - b)(x - c) \dots = 0.$$

5. Show that the coefficients of the equation are simple functions of the negatives of the roots.

6. Show that, if, on substituting two different values for x in turn in the first member of an equation, the results are of opposite signs, there must be an odd number of roots between the values substituted.

SOLUTION OF NUMERICAL EQUATIONS.

Commensurable Roots.

7. Prove that if the coefficients of an equation are whole numbers, and the coefficient of the first term is unity, the equation cannot have a fractional root. Any commensurable root of such an equation must then be an exact divisor of the constant term by § 5.

8. After a root a is found, the degree of the equation may be lowered by dividing by $x - a$. Explain abbreviated methods of division: first, by detached coefficients; second, by synthetic division.

9. Prove the theorem: If a is a commensurable root it will exactly divide the constant term, the quotient thus obtained plus the coefficient of the preceding term, this quotient plus the preceding coefficient, and so on, and the last quotient will be -1 .

Prove the converse of this theorem.

Give the working method of finding all the commensurable roots of the equation described in § 7.

10. If the coefficients of an equation are whole numbers, and the coefficient of the first term is not unity, show that the equation may easily be transformed into one where the coefficients are whole numbers and the coefficient of the first term is unity, and may then be treated by § 9.

11. *Descartes' Rule of Signs.* Explain what is meant by a *permanence* of sign; a *variation* of sign.

Prove that the number of positive roots of an equation, complete or incomplete, cannot exceed the number of variations of sign. Show that, by reversing the signs of the terms of odd degree, the equation may be transformed into one whose roots are the negatives of the roots of the given equation, and that by applying Descartes' Rule to the transformed equation, information may be obtained concerning the negative roots of the given equation.

Incommensurable Roots. Methods of Approximation.

12. Explain the rough laborious method of approximating to the value of an incommensurable root based upon § 6, and show that it is theoretically capable of any desired degree of accuracy, and as applicable to a transcendental as to an algebraic equation.

13. Explain briefly *Newton's Method*; *i.e.*, find a portion a of the root by § 12; let $x = a + h$, and substitute this value for x in the equation, neglecting higher powers of h than the first, thus obtaining an equation of the first degree to determine an approximate value of h . The result is reasonably accurate if h is small.

14. *Horner's Method.* Find by § 12 a portion a of the root. Transform the equation into one whose roots are less by a than those of the original equation. Treat the resulting equation by § 12.

Show that the coefficients of the transformed equation will be the remainders obtained in dividing the original equation repeatedly by $x - a$.

Newton's Method, § 13. shows that an approximate value for the rest of the root may be found by dividing the constant term of the transformed equation by the preceding coefficient.

Describe *Horner's Method* in its practical abbreviated working form. Show how to deal with negative roots.

General Methods and Theorems.

15. Explain the method of finding equal roots by obtaining the greatest common divisor of the first member of the equation and its derivative with respect to x .

16. Describe *Sturm's Functions*. Prove Sturm's Theorem.

17. In dealing with a given numerical equation of high degree: first, test for commensurable roots, and lower the degree of the equation by their aid if any are found; second, test for equal roots; third, use Horner's Method in finding approximately the incommensurable roots, employing Sturm's Theorem as an auxiliary if it proves absolutely necessary.

IMAGINARIES.

18. The treatment and use of *imaginaries* is purely arbitrary and conventional. Define the square root of -1 as a symbol of operation, and state the conventions adopted to govern its use.

$$\begin{aligned}\sqrt{-a^2} &= a\sqrt{-1}, \\ (a+b)\sqrt{-1} &= a\sqrt{-1} + b\sqrt{-1}, \\ a\sqrt{-1} &= \sqrt{-1}.a.\end{aligned}$$

Interpret the powers of $\sqrt{-1}$ by the aid of the definition and these conventions.

Show that these conventions enable us to deal with imaginary roots of a quadratic, and that their treatment and properties are closely analogous to those of real roots.

19. Show why $a + b\sqrt{-1}$ is taken as the typical form of an imaginary. Explain the ordinary geometrical representation of an imaginary by the position of a point in a plane. N.B. This interpretation is entirely arbitrary, but has proved very useful in suggesting important relations which might not otherwise have been discovered.

20. Show that the *sum*, the *product*, and the *quotient* of two imaginaries are imaginaries of the typical form.

21. Give the second typical form of an imaginary suggested by the graphical construction of § 19.

$$r(\cos \phi + \sqrt{-1} \cdot \sin \phi).$$

Define the *modulus* and the *argument* of an imaginary.

State the convention concerning the sign of the modulus, and show that the argument may have an infinite number of values differing by multiples of 2π .

22. Show that the modulus of the product of two imaginaries is the product of their moduli, and that the argument of their product is the sum of their arguments. Prove the theorems concerning the modulus and argument of the quotient of two imaginaries; of a power of an imaginary; of a root of an imaginary.

23. Show that the n th root of any real or imaginary has n values, having the same modulus and arguments differing by multiples of $\frac{2\pi}{n}$.

24. Define *conjugate* imaginaries. Prove that conjugate imaginaries have a real sum and a real product.

Show that if an equation with real coefficients has an imaginary root, the conjugate of that root is also a root of the equation.

25. Give *Cardan's Solution of a Cubic* of the form

$$x^3 + qx + r = 0.$$

Consider the *irreducible case*. Give a *Trigonometric Solution*. Show that any cubic can be reduced to the form

$$x^3 + qx + r = 0.$$

Obtain the general solution of any cubic.

26. Give *Descartes' and Euler's Methods* of solving a bi-quadratic equation.

SYMMETRIC FUNCTIONS OF THE ROOTS OF AN EQUATION.

27. Define a *symmetric function* of several quantities. Show that any combination of symmetric functions is symmetric.

The coefficients of an equation are symmetric functions of the roots of the equation by § 5.

28. Explain *Newton's Method* of expressing the sums of powers of the roots of an equation in terms of the coefficients,

$$fx = (x - a)(x - b)(x - c) \dots$$

Take the logarithm of each member and differentiate

$$\frac{f'x}{fx} = \frac{1}{x - a} + \frac{1}{x - b} + \frac{1}{x - c} \dots$$

Consider the case where the required power is less than the degree of the equation; where the required power is greater than the degree of the equation.

29. Give the short practical method of obtaining the sums of powers of the roots of a numerical equation; divide $xf'x$ by fx , and the coefficients of x^{-1} , x^{-2} , x^{-3} , etc., in the quotient are s_1 , s_2 , s_3 , etc. Shorten by using detached coefficients.

30. Any complicated symmetric function can be made to depend upon simpler functions so that only rational integral forms need be specially investigated.

31. Show that symmetric functions may be expressed in terms of the sums of powers of the quantities involved.

$$\Sigma a^m = s_m,$$

$$\Sigma a^m b^p = s_m s_p - s_{m+p},$$

$$\Sigma a^m b^p c^q = s_m s_p s_q - s_{m+p} s_q - s_{m+q} s_p - s_{p+q} s_m + 2s_{m+p+q}.$$

Consider special cases.

32. Explain the method of *elimination by the aid of symmetric functions*.

DETERMINANTS.

33. Show that, if two simultaneous equations of the first degree,

$$a_1 x + b_1 y + c_1 = 0,$$

$$a_2 x + b_2 y + c_2 = 0,$$

are solved, the numerators and denominators of the values of x and y have a peculiar symmetric form.

Explain the notation adopted for writing compactly such expressions.

Describe a *Determinant*, its *rows*, *columns*, and *diagonal term*.

Give a rule for expanding a determinant. Give the *law of signs*.

Illustrate by determinants of the second and third orders.

34. Show that a determinant may be broken up into a sum of terms each involving a *sub-determinant*. Illustrate.

35. Show that an interchange of two rows or of two columns will change the sign of a determinant.

36. Show that if two rows or two columns are identical, the value of the determinant is zero.

37. Show that if each constituent of any row or column is multiplied by a given quantity, the whole determinant is multiplied by that quantity.

38. Show that if each constituent of any row or of any column is a binomial, the determinant can be broken up into the sum of two other determinants of the same order.

39. Show how to compute the value of a numerical determinant. Consider examples.

40. Explain the application of determinants to *elimination* in the case of n equations of the first degree between n unknown quantities.

41. Explain the application of determinants to elimination in the case of two equations of any degree involving two unknown quantities.

W. E. BYERLY,

Professor of Mathematics in Harvard University.

ELEMENTS OF THE DIFFERENTIAL CALCULUS.

With Numerous Examples and Applications. Designed for Use as a College Text-Book. By W. E. BYERLY, Professor of Mathematics, Harvard University. 8vo. 273 pages. Mailing Price, \$2.15; Introduction, \$2.00.

This book embodies the results of the author's experience in teaching the Calculus at Cornell and Harvard Universities, and is intended for a text-book, and not for an exhaustive treatise. Its peculiarities are the rigorous use of the Doctrine of Limits, as a foundation of the subject, and as preliminary to the adoption of the more direct and practically convenient infinitesimal notation and nomenclature; the early introduction of a few simple formulas and methods for integrating; a rather elaborate treatment of the use of infinitesimals in pure geometry; and the attempt to excite and keep up the interest of the student by bringing in throughout the whole book, and not merely at the end, numerous applications to practical problems in geometry, and mechanics.

James Mills Peirce, Prof. of Math., Harvard University. From the *Harvard Register*: In mathematics, as in other branches of study, the need is now very much felt of teaching which is general without being superficial; limited to leading topics, and yet within its limits; thorough, accurate, and practical; adapted to the communication of some degree of power, as well as knowledge, but free from details which are important only to the specialist. Prof. Byerly's Calculus appears to be designed to meet this want. . . . Such a plan leaves much room for the exercise of individual judgment, and differences of opinion will undoubtedly exist in regard to one and another point of this book. But all teachers will agree that in selection, arrangement, and treatment, it is, on the whole, in a very high degree, wise, able, marked by a true scien-

tific spirit, and calculated to develop the same spirit in the learner. . . . The book contains, perhaps, all of the integral calculus, as well as of the differential, that is necessary to the ordinary student. And with so much of this great scientific method, every thorough student of physics, and every general scholar who feels any interest in the relations of abstract thought, and is capable of grasping a mathematical idea, ought to be familiar. One who aspires to technical learning must supplement his mastery of the elements by the study of the comprehensive theoretical treatises. . . . But he who is thoroughly acquainted with the book before us has made a long stride into a sound and practical knowledge of the subject of the calculus. He has begun to be a real analyst.

ELEMENTS OF THE INTEGRAL CALCULUS.

With Numerous Examples and Applications, containing a Chapter on the Calculus of Imaginaries, and a Practical Key to the Solution of Differential Equations. Designed for Use as a College Text-Book. By W. E. BYERLY, Prof. of Mathematics in Harvard University. 8vo. 204 pages. Mailing Price, \$2.15. Introduction, \$2.00.

This volume is a sequel to the author's treatise on the Differential Calculus, and, like that, is written as a text-book. The last chapter, however, — a Key to the Solution of Differential Equations, — may prove of service to working mathematicians.

H. A. Newton, Professor of Mathematics, Yale College: We shall use it in my optional class next term.

UNIVERSITY OF CALIFORNIA LIBRARY
BERKELEY

Return to desk from which borrowed.
This book is DUE on the last date stamped below.

MAR 25 1948

JUL 15 1950 R

~~_____~~

~~_____~~
~~_____~~

8 Nov '57 R K

OCT 1 1971 8 8

RECD LD 801 1 71-9AM 6 8

GAYLAMOUNT
PAMPHLET BINDER



Manufactured by
GAYLORD BROS. Inc.
Syracuse, N. Y.
Stockton, Calif.

U C BERKELEY LIBRARIES



061348240

M521572

